
DNA MICROMAVE RESPONSE(U) PURDUE UNIV LAFAVETTE IND DEPT OF PHYSICS L L VAN ZANDT 30 APR 87 UNCLASSIFIED F/G 28/14 NL

1/1

AD-A189 253



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A

AD-A180 25



20. DISTRIBUTION/AVAILABILITY OF ABSTRACT	21. ABSTRACT SECURITY CLASSIFICATION	
₩ UNCLASSIFIED/UNLIMITED	(U)	
22a NAME OF RESPONSIBLE INDIVIDUAL	22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL
M. Marron	202/696-4038	ONR

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

Response of DNA in Aqueous Solution to Microwave Irradiation

M.E. Davis & L.L. Van Zandt
 Department of Physics
 Purdue University
 W. Lafayette, Indiana 47907

We have calculated the expected absorption of microwave radiation in the gigaHertz frequency range by fixed length DNA polymer molecules disolved in saline solution. the effects of counterions and solvent dynamics have been accounted for in detail, the features of the absorption are completelly dominated by the interaction between charged polymer molecule and the so-called first hydration that is, the nearest layer of solvent water layer, molecules not actually bonded to the polymer. The relevant parameters of this interaction are the strength of the water-to-polymer coupling and the average persistence time of individual water-to-polymer bonds. These are presumably bonds to the oxygen atoms of the backbone hydrogen phosphate structure. Using the best available estimates of the system parameters, we can obtain structured absorption corresponding to compressional wave phonon excitations on the polymer, "organ pipe" modes, such as have been claimed to be seen by Edwards, Davis, Swicord & Saffer. These modes are now well known in much drier sample materials. We have examined the effects of varying solvent salt concentration and the system temperature. In both cases, the effects are virtually nil, in the former because of the Manning condensation phenomenon that preserves a remarkably constant polymer environment over a wide range of bulk and in the later case because of a ionic strength, fortuitous competition between effects of bulk viscosity and persistence time changes with temperature. Hence, any effects seen in the experimental variation of temperature salinity could be wholly attributed to their modification of the hydration layer properties.

The Effect of Propeller Twist on Phonon Dispersion in DNA

L. L. VanZandt and W. K. Schroll

Department of Physics
Purdue University
West Lafayette, Indiana 47907

We have developed a high accuracy software package for calculating a complete phonon dispersion spectrum for any DNA polymer configuration utilizing classical dynamics via a dynamical force constant matrix. These routines permit us to treat a variety of related physical systems with a minimum of attention to computing detail. As an example we have varied selected mean helix parameters over their allowed experimental ranges and then determined the particular values which yield a Helmholts free energy minimum. A purine-pyrimidine base pair has a propeller twist angle defined as the angular displacement between the planes of the two bases when they are rotated in opposite directions about a common axis determined by the nitrogen-carbon bond joining each base with its associated sugar. This paper deals with the optimization of the propeller twist angle, & for B-configuration Poly(dG) Poly(dC) DNA, which has an X-ray experimental uncertainty of ±4.8°. We show a marked variation in phonon frequencies at zone center for the lower optical modes over the 9.6° variance permitted experimentally. This is to be expected because the two acoustic modes are mediated by the bulk structure of the polymer rather than the particular atomic positions. The highest optical modes are unaffected since these high frequency phonons generally correspond to oscillations of relatively localized atomic coordinates. We evaluate these various configurations and determine that a minimum in the Helmholtz free energy corresponds to a propeller twist angle 2.20° less than that quoted as the equilibrium value. Further applications of this algorithm package for DNA structure parameter determination will include baseroll, base inclination and B to Z conformational change.